

# Contents

1	Python Classes		3
	1.1 Mak	king a Polynomial class	3
2	Common Polynomial Products		9
	2.1 Diffe	ference of squares	9
	2.2 Pow	vers of binomials	12
3	Factoring Polynomials		15
	3.1 How	w to factor polynomials	16
4	Partial Fractions		19
	4.0.	.1 Improper fractions	20
	4.0.2	.2 Proper fractions	21
5	Practice with Polynomials		
A	Answers to Exercises		
Index			31

CHAPTER 1

# Python Classes

The built-in types, such as strings, have functions associated with them. So, for example, if you needed a string converted to uppercase, you would call its upper() function: -

```
my_string = "houston, we have a problem!"
louder_string = my_string.upper()
```

This would set louder\_string to "HOUSTON, WE HAVE A PROBLEM!" When a function is associated with a datatype like this, it called a *method*. A datatype with methods is known as a *class*. The data of that type is known as the *instance* of that class. For example, in the example, we would say "my\_string is an instance of the class str. str has a method called upper"

The function type will tell you the type of any data:

```
print(type(my_string))
```

This will output

```
<class 'str'>
```

A class can also define operators. +, for example, is redefined by str to concatenate strings together:

long\_string = "I saw " + "15 people"

### 1.1 Making a Polynomial class

You have created a bunch of useful python functions for dealing with polynomials. Notice how each one has the word "polynomial" in the function name like derivative\_of\_polynomial. Wouldn't it be more elegant if you had a Polynomial class with a derivative method? Then you could use your polynomial like this:

```
a = Polynomial([9.0, 0.0, 2.3])
b = Polynomial([-2.0, 4.5, 0.0, 2.1])
```

```
print(a, "plus", b , "is", a+b)
print(a, "times", b , "is", a*b)
print(a, "times", 3 , "is", a*3)
print(a, "minus", b , "is", a-b)
c = b.derivative()
print("Derivative of", b ,"is", c)
And it would output:
2.30x^2 + 9.00 plus 2.10x^3 + 4.50x + -2.00 is 2.10x^3 + 2.30x^2 + 4.50x + 7.00
2.30x^2 + 9.00 times 2.10x^3 + 4.50x + -2.00 is 4.83x^5 + 29.25x^3 + -4.60x^2 + 40.50x + -18.00
2.30x^2 + 9.00 times 3 is 6.90x^2 + 27.00
2.30x^2 + 9.00 minus 2.10x^3 + 4.50x + -2.00 is -2.10x^3 + 2.30x^2 + -4.50x + 11.00
Derivative of 2.10x^3 + 4.50x + -2.00 is 6.30x^2 + 4.50
```

Create a file for your class definition called Polynomial.py. Enter the following:

```
class Polynomial:
    def __init__(self, coeffs):
        self.coefficients = coeffs.copy()
    def __repr__(self):
        # Make a list of the monomial strings
        monomial_strings = []
        # For standard form we start at the largest degree
        degree = len(self.coefficients) - 1
        # Go through the list backwards
        while degree >= 0:
            coefficient = self.coefficients[degree]
            if coefficient != 0.0:
                # Describe the monomial
                if degree == 0:
                    monomial_string = "{:.2f}".format(coefficient)
                elif degree == 1:
                    monomial_string = "{:.2f}x".format(coefficient)
                else:
                    monomial_string = "{:.2f}x^{}".format(coefficient, degree)
                # Add it to the list
                monomial_strings.append(monomial_string)
```

```
# Move to the previous term
        degree = degree - 1
    # Deal with the zero polynomial
    if len(monomial_strings) == 0:
        monomial_strings.append("0.0")
    # Separate the terms with a plus sign
    return " + ".join(monomial_strings)
def __call__(self, x):
   sum = 0.0
    for degree, coefficient in enumerate(self.coefficients):
        sum = sum + coefficient * x ** degree
    return sum
def __add__(self, b):
    result_length = max(len(self.coefficients), len(b.coefficients))
   result = []
    for i in range(result_length):
        if i < len(self.coefficients):</pre>
            coefficient_a = self.coefficients[i]
        else:
            coefficient_a = 0.0
        if i < len(b.coefficients):</pre>
            coefficient_b = b.coefficients[i]
        else:
            coefficient_b = 0.0
        result.append(coefficient_a + coefficient_b)
    return Polynomial(result)
def __mul__(self, other):
    # Not a polynomial?
    if not isinstance(other, Polynomial):
        # Try to make it a constant polynomial
        other = Polynomial([other])
    # What is the degree of the resulting polynomial?
    result_degree = (len(self.coefficients) - 1) + (len(other.coefficients) - 1)
    # Make a list of zeros to hold the coefficents
    result = [0.0] * (result_degree + 1)
```

```
# Iterate over the indices and values of a
    for a_degree, a_coefficient in enumerate(self.coefficients):
        # Iterate over the indices and values of b
        for b_degree, b_coefficient in enumerate(other.coefficients):
            # Calculate the resulting monomial
            coefficient = a_coefficient * b_coefficient
            degree = a_degree + b_degree
            # Add it to the right bucket
            result[degree] = result[degree] + coefficient
    return Polynomial(result)
__rmul__ = __mul__
def __sub__(self, other):
    return self + other * -1.0
def derivative(self):
    # What is the degree of the resulting polynomial?
    original_degree = len(self.coefficients) - 1
    if original_degree > 0:
        degree_of_derivative = original_degree - 1
    else:
        degree_of_derivative = 0
    # We can ignore the constant term (skip the first coefficient)
    current degree = 1
    result = []
    # Differentiate each monomial
    while current_degree < len(self.coefficients):
        coefficient = self.coefficients[current_degree]
        result.append(coefficient * current_degree)
        current_degree = current_degree + 1
    # No terms? Make it the zero polynomial
    if len(result) == 0:
        result.append(0.0)
    return Polynomial(result)
```

Create a second file called test\_polynomial.py to test it:

```
from Polynomial import Polynomial
a = Polynomial([9.0, 0.0, 2.3])
b = Polynomial([-2.0, 4.5, 0.0, 2.1])
print(a, "plus", b , "is", a+b)
print(a, "times", b , "is", a*b)
print(a, "times", 3 , "is", a*3)
print(a, "minus", b , "is", a-b)
c = b.derivative()
print("Derivative of", b ,"is", c)
slope = c(3)
print("Value of the derivative at 3 is", slope)
```

Run the test code:

python3 test\_polynomial.py

## CHAPTER 2

# **Common Polynomial Products**

In math and physics, you will run into certain kinds of polynomials over and over again. In this chapter, We are going to cover some patterns that you will want to be able to recognize.

### 2.1 Difference of squares

Watch **Polynomial special products: difference of squares** from Khan Academy at https://youtu.be/uNweU6I4Icw.

If you are asked what (3x-7)(3x+7) is, you would use the distributive property to expand that to (3x)(3x) + (3x)(7) + (-7)(3x) + (-7)(7). Two of the terms cancel each other, so this is  $(3x)^2 - (7)^2$ . This would simplify to  $9x^2 - 49$ 

You will see this pattern often. Anytime you see (a + b)(a - b), you should immediately recognize it equals  $a^2-b^2$ . (Note that the order doesn't matter: (a-b)(a+b) also  $a^2-b^2$ .)

Working the other way is important too. Any time you see  $a^2 - b^2$ , that you should recognize that you can change that into the product (a + b)(a - b). Making something into a product like this is known as *factoring*. You probably have done prime factorization of numbers like  $42 = 2 \times 3 \times 7$ . In the next couple of chapters, you will learn to factorize polynomials.

### **Exercise 1** Difference of Squares



We are often interested in the roots of a polynomial. That is, we want to know "For what values of x does the polynomial evaluate to zer?" For example, when you deal with falling bodies, the first question you might ask would be "How many seconds before the hammer hits the ground?" Once you have factored a polynomial into binomials, you can easily find the roots.

For example, what are the roots of  $x^2 - 5$ ? You just factored it into  $(x + \sqrt{5})(x - \sqrt{5})$  This product is zero if and only if one of the factors is zero. The first factor is only zero when x is  $-\sqrt{5}$ . The second factor is zero only when x is  $\sqrt{5}$ . Those are the only two roots of this polynomial.

Let's check that result.  $\sqrt{5}$  is a little more than 2.2. Using your Python code, you can graph the polynomial:

import poly.py
import matplotlib.pyplot as plt
# x\*\*2 - 5
pn = [-5.0, 0.0, 1.0]

```
# These lists will hold our x and y values
x_list = []
y_list = []
# Start at x=-3
current_x = -3.0
# End at x=3.0
while current_x < 3.0:
    current_y = poly.evaluate_polynomial(pn, current_x)
    # Add x and y to respective lists
    x_list.append(current_x)
    y_list.append(current_y)
    # Move x forward
    current_x += 0.1
# Plot the curve
plt.plot(x_list, y_list)
plt.grid(True)
plt.show()
```

You should get a plot like this:



It does, indeed, seem to cross the x-axis near -2.2 and 2.2.

### 2.2 **Powers of binomials**

You can raise whole polynomials to exponents. For example,

$$(3x^3 + 5)^2 = (3x^3 + 5)(3x^3 + 5)$$
  
= 9x<sup>6</sup> + 15x<sup>3</sup> + 15x<sup>3</sup> + 25 = 9x<sup>6</sup> + 30x<sup>3</sup> + 25

A polynomial with two terms is called a *binomial*.  $5x^9 - 2x^4$ , for example, is a binomial. In this section, we are going to develop some handy techniques for raising a binomial to some power.

Looking at the previous example, you can see that for any monomials a and b,  $(a+b)^2 = a^2 + 2ab + b^2$ . So, for example,  $(7x^3 + \pi)^2 = 49x^6 + 14\pi x^3 + \pi^2$ 

Working Space

## **Exercise 2** Squaring binomials

Simply the following

- 1.  $(x+1)^2$
- 2.  $(3x^5+5)^2$
- 3.  $(x^3 1)^2$
- 4.  $(x \sqrt{7})^2$

\_\_\_\_\_ Answer on Page 29 \_\_\_\_\_

What about  $(x + 2)^3$ ? You can do it as two separate multiplications:

$$(x+2)^3 = (x+2)(x+2)(x+2)$$
  
= (x+2)(x<sup>2</sup>+4x+4) = x<sup>3</sup>+4x<sup>2</sup>+4x+2x<sup>2</sup>+8x+8  
= x<sup>3</sup>+6x<sup>2</sup>+12x+8

In general, we can say that for any monomials a and b,  $(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$ .

What about higher powers?  $(a+b)^4$ , for example? You could use the distributive property four times, but it starts to get pretty tedious.

Here is a trick. This is known as *Pascal's triangle* 

Each entry is the sum of the two above it.

The coefficients of each term are given by the entries in Pascal's triangle:

$$(a+b)^4 = 1a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + 1b^4$$

## **Exercise 3** Using Pascal's Triangle

 1. What is  $(x + \pi)^5$ ?

 Answer on Page 30

## CHAPTER 3

# **Factoring Polynomials**

We factor a polynomial into two or more polynomials of lower degree. For example, let's say that you wanted to factor  $5x^3 - 45x$ . You would note that you can factor out 5x from every term. Thus,

$$5x^3 - 45x = (5x)(x^2 - 9)$$

You might notice that the second factor looks like the difference of squares, so

 $5x^3 - 45x = (5x)(x+3)(x-3)$ 

That is as far as we can factorize this polynomial.

Why do we care? The factors make it easy to find the roots of the polynomial. This polynomial evaluates to zero if and only if at least one of the factors is zero. Here, we see that

- The factor (5x) is zero when x is zero.
- The factor (x + 3) is zero when x is -3.
- The factor(x 3) is zero when x is 3.

So, looking at the factorization, you can see that  $5x^3 - 45x$  is zero when x is 0, -3, or 3.

This is a graph of that polynomial with its roots circled:



### 3.1 How to factor polynomials

The first step when you are trying to factor a polynomial is to find the greatest common divisor for all the terms, then pull that out. In this case, the greatest common divisor will also be a monomial. Its degree is the least of the degrees of the terms, its coefficient will be the greatest common divisor of the coefficients of the terms.

For example, what can you pull out of this polynomial?

$$12x^{1}00 + 30x^{3}1 + 42x^{1}7$$

The greatest common divisor of the coefficients (12, 30, and 42) is 6. The least of the degrees of terms (100, 31, and 17) is 17. So, you can pull out  $6x^{1}7$ :

$$12x^{1}00 + 30x^{3}1 + 42x^{1}7 = (6x^{1}7)(2x^{8}3 + 5x^{1}4 + 7)$$

### **Exercise 4** Factoring out the GCD monomial



So, now you have the product of a monomial and a polynomial. If you are lucky, the polynomial part looks familiar, like the difference of squares or a row from Pascal's triangle.

Often, you are trying factor a quadratic like  $x^2 + 5x + 6$  in a pair of binomials. In this case, the result would be (x + 3)(x + 2). Let's check that:

 $(x+3)(x+2) = (x)(x) + (3)(x) + (2)(x) + (3)(2) = x^2 + 5x + 6$ 

Notice that 3 and 2 multiply to 6 and add to 5. If you were trying to factor  $x^2 + 5x + 6$ , you would ask yourself" What are two numbers that when multiplied equal 6 and when added equal 5?" And you might guess wrong a couple of times. For example, you might say to youself, "Well, 6 times 1 is 6. Maybe those work. But 6 and 1 add 7. So those don't work."

Solving these sorts of problems are like solving a Sudoku puzzle. You try things and realize they are wrong, so you backtrack and try something else.

The numbers are sometimes negative. For example,  $x^2 + 3x - 10$  factors into (x+5)(x-2).

## **Exercise 5 Factoring quadratics**



## CHAPTER 4

## **Partial Fractions**

How can you add fractions with different denominators, like  $\frac{1}{x} + \frac{2}{x+3}$ ? You would need to make the denominators the same; after that, you could just add the numerators. You achieve this by multiplying the numerator and denominator of each fraction by the denominator of the other fraction:

$$\frac{1}{x} + \frac{2}{x+3} = \frac{1}{x} \left( \frac{x+3}{x+3} \right) + \frac{2}{x+3} \left( \frac{x}{x} \right)$$

Recall that when the numerator and denominator of a fraction are the same, the fraction is equal to one. So, we are not changing the *value* of each fraction, since we are just multiplying by one. Continuing, we can perform the multiplication and see that:

$$\frac{1}{x} + \frac{2}{x+3} = \frac{x+3}{x(x+3)} + \frac{2x}{x(x+3)}$$
$$= \frac{(x+3)+2x}{x(x+3)} = \frac{3x+3}{x^2+3x} = \frac{3(x+1)}{x^2+3x}$$

The inverse of this process is called **partial fraction decomposition** (or partial fraction expansion). This method has applications in many fields, but we will find it most useful as a tool to evaluate integrals in a later chapter.

Let g(x) be a rational function such that

$$g(\mathbf{x}) = \frac{\mathsf{P}(\mathbf{x})}{\mathsf{Q}(\mathbf{x})}$$

Where P(x) and Q(x) are polynomials. If g(x) is proper (that is, the degree of P is less than the degree of Q) then we can express g(x) as the sum of simpler rational fractions. If g(x) is improper (that is, the degree of P is greater than or equal to the degree of Q), then we must first perform long division to obtain a remainder, R(x), where the degree of R is less than the degree of Q:

$$g(x) = \frac{P(x)}{Q(x)} = S(x) + \frac{R(x)}{Q(x)}$$

#### 4.0.1 Improper fractions

What is  $\int \frac{x^3 + x}{x - 1} dx$ . Using long division, we see that:

$$\frac{x^3 + x}{x - 1} = x^2 + x + 2 + \frac{2}{x - 1}$$

(see figure 4.1 for an explanation). Then we can also say that:

$$\frac{x^{3} + x}{x - 1} = x^{2} + x + 2 + \frac{2}{x - 1}$$

$$x - 1 \left[ \begin{array}{c} x^{2} + x + 2 \\ x - 1 \end{array} \right] \left[ \begin{array}{c} x^{2} + x + 2 \\ x^{3} + 0x^{2} + x \\ -(x^{3} - x^{2}) \\ x^{2} + x \\ -(x^{2} - x) \\ 2x \\ -(2x - 2) \\ 2 \end{array} \right]$$

Figure 4.1: Evaluating  $(x^3 + x) \div (x - 1)$  with the long division method

When you start with an improper fraction, use long division to reduce it to a term plus a proper fraction, then use the methods outlined below to further manipulate the proper fraction.

#### **Exercise 6**

Use long division to reduce the following improper rational functions to a term plus a proper rational fraction.

1.  $\frac{x^4 + x^3 + 2x^2 + 2x - 3}{x^2 - 3x + 2}$ 2.  $\frac{2x^3 + 5}{x^3 - 3x^2 + 2x - 4}$ 

3. 
$$\frac{3x^4 - 2x^3 - x^2 + 1}{x^3 - 3x}$$

Working Space

Answer on Page 30

#### 4.0.2 **Proper fractions**

When the order of the numerator is less than or equal to the denominator, there are three further possibilities.

#### No repeated linear factors

In the first case, the denominator, Q(x) is composed of distinct linear factors. In this case, we can say that  $Q(x) = (a_1x + b_1)(a_2x + b_2) \cdots (a_nx + b_n)$ , where no factor is repeated (including constant multiples). Then, there exists A, B, C,  $\cdots$ , such that:

$$\frac{P(x)}{Q(x)} = \frac{A}{a_1x + b_1} + \frac{B}{a_2x + b_2} + \cdots$$

Let's see an example of this by decomposing  $\frac{4x^2-7x-12}{x(x+2)(x-3)}$ . We start by defining A, B, and C, such that:

$$\frac{4x^2 - 7x - 12}{x(x+2)(x-3)} = \frac{A}{x} + \frac{B}{x+2} + \frac{C}{x-3}$$

Multiplying both sides by x(x+2)(x-3) we get:

$$4x^2 - 7x - 12 = A(x+2)(x-3) + B(x)(x-3) + C(x)(x+2)$$

We have 3 unknowns and only one equation! Don't worry — remember this equation is true for all x, so we can choose a convenient value of x to isolate each unknown in turn. Starting, let x = 0. Then:

$$4(0)^{2} - 7(0) - 12 = A(0+2)(0-3) + B(0)(x-3) + C(0)(x+2)$$
$$-12 = A(2)(-3) + 0 + 0$$

Notice that the B and C disappear, and we can solve for A:

$$A = \frac{-12}{-6} = 2$$

We can solve for B by setting x = -2 and for C by setting x = 3 (notice, we've used all three zeroes of the denominator polynomial):

$$4(-2)^{2} - 7(-2) - 12 = A(-2+2)(-2-3) + B(-2)(-2-3) + C(-2)(-2+2)$$

$$4(4) + 14 - 12 = 0 + B(-2)(-5) + 0$$

$$16 + 2 = 10B$$

$$B = \frac{9}{5}$$

and

$$4(3)^{2} - 7(3) - 12 = A(3+2)(3-3) + B(3)(3-3) + C(3)(3+2)$$
  

$$4(9) - 21 - 12 = 0 + 0 + C(3)(5)$$
  

$$36 - 33 = 15C$$
  

$$C = \frac{1}{5}$$

We can then decompose our original fraction:

$$\frac{4x^2 - 7x - 12}{x(x+2)(x-3)} = \frac{2}{x} + \frac{9}{5(x+2)} + \frac{1}{5(x-3)}$$

You can check your answer by cross-multiplying and adding. You should get the same rational function back.

#### **Repeated linear factors**

The second case is if Q(x) has repeated factors (such as  $x^2 + 8x + 16 = (x + 4)^2$ ). Suppose the first linear factor,  $(a_1x + b_1)$  is repeated r times (that is, Q(x) contains the factor  $(a_1x + b_1)^r$ ). Then, instead of  $\frac{A}{a_1x+b_1}$ , we should write:

$$\frac{A_1}{a_1x + b_1} + \frac{A_2}{(a_1x + b_1)^2} + \dots + \frac{A_r}{(a_1x + b_1)^r}$$

Let's look at a concrete example to see how this works:

**Example:** Decompose  $\frac{x^2+x+1}{(x+1)^2(x+2)}$ 

**Solution**: We start by defining:

$$\frac{x^2 + x + 1}{(x+1)^2(x+2)} = \frac{A}{x+1} + \frac{B}{(x+1)^2} + \frac{C}{x+2}$$

Multiplying both sides by  $(x + 1)^2(x + 2)$ :

$$x^{2} + x + 1 = A(x + 1)(x + 2) + B(x + 2) + C(x + 1)^{2}$$

Since there are only 2 roots to  $(x+1)^2(x+2)$ , we will use another method called "equating the coefficients" to find A, B, and C. We start by expanding the right side of the equation:

$$x^{2} + x + 1 = A(x^{2} + 3x + 2) + B(x + 2) + C(x^{2} + 2x + 1)$$

Distributing and combining, we find that:

$$x^{2} + x + 1 = Ax^{2} + 3Ax + 2A + Bx + 2B + Cx^{2} + 2Cx + C$$

$$x^{2} + x + 1 = (A + C)x^{2} + (3A + B + 2C)x + (2A + 2B + C)$$

For this equation to be true, we know that:

$$A + C = 1$$
$$3A + B + 2C = 1$$
$$2A + 2B + C = 1$$

(That is, the coefficient for  $x^2$  on the left, 1, must be equal to the coefficient for  $x^2$  on the right, (A + C), and so on.) We now have a system of 3 equations and 3 unknowns. When you solve for each, you should find that:

$$A = -2$$
$$B = 1$$
$$C = 3$$

Therefore,

$$\frac{x^2 + x + 1}{(x+1)^2(x+2)} = \frac{-2}{x+1} + \frac{1}{(x+1)^2} + \frac{3}{x+2}$$

#### Irreducible quadratic factors

Sometimes, we cannot express a polynomial as the product of two linear statements (that is, terms in the form ax + b). Take  $x^2 + 1$ , which cannot be expressed as the product of real, linear terms. What do you do if something like  $x^2 + 1$  is in the denominator? In this case, when we write an expression for  $\frac{P(x)}{Q(x)}$ , we include a term in the form:

$$\frac{Ax + B}{ax^2 + bx + c}$$

For example, we can write:

$$\frac{x}{(x-2)(x^2+1)(x^2+4)} = \frac{A}{x-2} + \frac{Bx+C}{x^2+1} + \frac{Dx+E}{x^2+4}$$

**Example:** Decompose  $\frac{2x^2-x+4}{x^3+4x}$ 

Solution: We begin by factoring the denominator:

$$x^3 + 4x = x(x^2 + 4)$$

Which cannot be factored further. Therefore, we define:

$$\frac{2x^2 - x + 4}{x(x^2 + 4)} = \frac{A}{x} + \frac{Bx + C}{x^2 + 4}$$
$$2x^2 - x + 4 = A(x^2 + 4) + (Bx + C)x$$
$$2x^2 - x + 4 = Ax^2 + 4A + Bx^2 + Cx$$

Which implies that:

$$2 = A + B$$
$$C = -1$$
$$4A = 4$$

Therefore, A = 1, B = 1, and C = -1 and we can say that:

$$\frac{2x^2 - x + 4}{x^3 + 4x} = \frac{1}{x} + \frac{x - 1}{x^2 + 4}$$

#### Repeated irreducible quadratic factors

Lastly, the denominator might contain repeated irreducible quadratic factors. Similar to repeated linear factors, when setting up your partial fractions, instead of only writing

$$\frac{A}{ax^2 + bx + c}$$

For a quadratic factor that is repeated r times, your equation should include:

$$\frac{A_1}{ax^2 + bx + c} + \frac{A_2}{(ax^2 + bx + c)^2} + \dots + \frac{A_r}{(ax^2 + bx + c)^r}$$

### Exercise 7

Decompose the following proper fractions

- 1.  $\frac{x-4}{x^2+5x-6}$
- 2.  $\frac{x^2 + x + 1}{(x^2 + 1)^2}$
- 3.  $\frac{x^2+x+1}{(x+1)^2(x+2)}$

\_\_\_\_\_ Answer on Page 30 \_\_\_\_

– Working Space

# Practice with Polynomials

At this point, you know all the pieces necessary to solve problems involving polynomials. In this chapter, you are going to practice using all of these ideas together.

Watch Khan Academy's **Polynomial identities introduction here**: https://youtu.be/ EvNKKyhLSpQ Also, watch the follow up here: https://youtu.be/-6qi049Q180

FIXME: Lots of practice problems here

## APPENDIX A

# Answers to Exercises

## Answer to Exercise 1 (on page 10)

$$(2x - 3)(2x + 3) = 4x^{2} - 9$$

$$(7 + 5x^{3})(7 - 5x^{3}) = 49 - 25x^{6}$$

$$(x - a)(x + a) = x^{2} - a^{2}$$

$$(3 - \pi)(3 + \pi) = 9 - \pi^{2}$$

$$(-4x^{3} + 10)(-4x^{3} - 10) = 16x^{6} - 100$$

$$(x + \sqrt{7})(x - \sqrt{7}) = x^{2} - 7$$

$$x^{2} - 9 = (x + 3)(x - 3)$$

$$49 - 16x^{6} = (7 + 4x^{3})(7 + 4^{3})$$

$$\pi^{2} - 25x^{8} = (\pi + 5x^{4})(\pi - 5x^{4})$$

$$x^{2} - 5 = (x + \sqrt{5})(x - \sqrt{5})$$

## Answer to Exercise 2 (on page 13)

$$(x + 1)^{2} = x^{2} + 2x + 1$$
  
$$(3x^{5} + 5)^{2} = 9x^{1}0 + 30x^{5} + 25$$
  
$$(x^{3} - 1)^{2} = x^{6} - 2x^{3} + 1$$
  
$$(x - \sqrt{7})^{2} = x^{2} - 2x\sqrt{7} + 7$$

## Answer to Exercise 3 (on page 14)

 $(x+\pi)^5 = x^5 + 5\pi x^4 + 10\pi^2 x^3 + 10\pi^3 + x^2 + 5\pi^2 x + \pi^5$ 

Answer to Exercise 4 (on page 17)

## Answer to Exercise 5 (on page 17)

## Answer to Exercise 6 (on page 20)

1. 
$$x^2 + 4x + 12\frac{30x-27}{x^2-3x+2}$$

2. 
$$2 + \frac{6x^2 - 4x + 13}{x^3 - 3x^2 + 2x - 4}$$

3. 
$$3x - 2 + \frac{8x^2 - 6x + 1}{x^3 - 3x}$$

## Answer to Exercise 7 (on page 25)

1. 
$$\frac{10}{7(x+6)} + \frac{-3}{7(x-1)}$$

2. 
$$\frac{1}{x^2+1} + \frac{x}{(x^2+1)^2}$$

3. 
$$\frac{-2}{x+1} + \frac{1}{(x+1)^2} + \frac{3}{x+2}$$



# INDEX

class in python, 3 factoring polynomials, 15 partial fraction decomposition, 19